

**RESEARCH ARTICLE** 

# The Effect Of Sea Cucumber Powder And Hyperbaric Oxygen Therapy On The Expression Of Tumor Necrosis Factor Alfa In Rats With Diabetes Mellitus Induced By Porphyromonas gingivalis

Caroline Prajnaparamitha Anggraini\*, Kristanti Parisihni\*\*, Widyastuti\*\*\*

\*Undergraduate Dentistry Faculty Hang Tuah University Surabaya

\*\*\*Oral Biology Faculty Hang Tuah University Surabaya

\*\*\*\*Periodontia Faculty Hang Tuah University Surabaya

#### **ABSTRACT**

Background: Periodontitis have a bidirectional relationship with diabetes mellitus, both can increase oxidative stress that trigger an increase of pro-inflammatory cytokines, such as TNF-α. Sea cucumbers have anti-inflammatory component that act to inhibit the release of inflammatory mediators. Hyperbaric oxygen therapy reduce oxidative stress. Purpose: The aim of this study is to investigate the effect of the sea cucumber powder and hyperbaric oxygen therapy on the expression of TNF-a in rats with diabetes mellitus induced by P.gingivalis bacteria. Materials and Methods: The study was an experimental laboratories research with factorial design. Twenty wistar rats were divided into 5 groups. K0 negative control, K1 positive control. K1-K4 groups were induced with streptozotocin for diabetes condition and P.gingivalis bacteria for periodontitis condition. K2 group was treated with sea cucumber 3% for 7 days, K3 with OHB 2.4 ATA 3 x 30' interval 5' for 7 days, and K4 group was treated with combination of sea cucumber and OHB. At the 51st day all rats were sacrified, then the expression of TNF-α on periodontal macrophages were examined by immunohistochemistry stain. Data were analized by Kruskal Wallis followed by Mannwhitney. **Result:** Expression of TNF- $\alpha$  were increased in K1 (11.50 $\pm$ 1.291) compare K0 (p<0.05). Sea cucumber treatment, OHB, and combination treatment decreased expression of TNF- $\alpha$  significantly in amount of 2.50±0.577 (K2), 8.25±2.217 (K3), 3.00±0.816 (K4). **Conclusion:** Sea cucumber powder and OHB therapy affected the expression of TNF- $\alpha$  in rats with diabetes mellitus induced by P. gingivalis bacteria.

**Keyword:** Sea cucumber powder, TNF- $\alpha$ , periodontitis, diabetes mellitus, OHB therapy

Correspondence: Kristanti Parisihni, Department Oral Biology, Faculty of Dentistry, Hang Tuah University, Arif Rahman Hakim 150, Surabaya, Phone 031-5945864, Email: <a href="mailto:tanti\_kris@yahoo.co.id">tanti\_kris@yahoo.co.id</a>



# BACKGROUND

The relationship between periodontitis and diabetes mellitus has long been discussed, both diseases have a relatively high incidence. Diabetes is a risk factor which increasing the severity of periodontitis. Instead, periodontitis may be a risk factor for glycemic control in people with diabetes.<sup>1,2</sup> Approximately 177 million people worldwide have diabetes mellitus and diabetes has been confirmed as a major risk factor for periodontitis. The risk of periodontitis increased approximately three-fold in individuals with diabetes with non-diabetic compared individuals.3,4

Individuals with diabetes have high glucose levels in the gingival fluid and blood more than individuals without diabetes. Increased glucose levels in the gingival fluid and blood in individuals with diabetes can change microflora environment, induce qualitative changes of bacteria that contribute to the severity, periodontal disease in patients uncontrolled diabetes. glycemic control led to increasing of oxidative stress, advanced glycation end products (AGEs), hyperresponsive innate immune system that contribute to damage factor diabetic periodontitis. In addition, the function of the normal immune cells, such as chemotaxis, cell phagocytosis, adherence, cytokine production and secretion, are all affected by hyperglycemia.<sup>5,6</sup>

Diabetics have more severe periodontal disease because of the presence of glucose which induce accumulation of advanced glycation end products (AGEs) and affect the activity of migration and phagocytic activity of mononuclear and polymorphonuclear, so the formation of a more pathogenic subgingival flora.<sup>7</sup> Many bacteria associated with periodontal disease including Porphyromonas gingivalis one of the most important periodontal pathogens. At the time of periodontal infection, the balance of bacterial flora in the oral cavity become changed, so the species of pathogenic bacteria is grow and survive, even when confronted with response.6 the immune Lipopolysaccharide (LPS) is one of most important microbial virulence factors in generating a host-mediated periodontal tissue destruction than that, lipopolysaccharide is a potent stimulator of various inflammatory mediators, including IL-1β and TNF-α.8

ISSN: 1907-5987

Tumor necrosis factor (TNF  $\alpha$ ) is reported have a role as a key to the pathogenesis of diabetes mellitus type 2. Research in animals and in vitro human, TNF- $\alpha$  inhibits the action of insulin. In animal and human cell models in vitro, TNF- $\alpha$  inhibits insulin. In one study, an increase in circulating levels of TNF- $\alpha$  is associated with insulin resistance and type 2 diabetes. Individuals with periodontitis disease have high TNF- $\alpha$  response, monocytes from individuals with diabetes and periodontitis release massive amounts of TNF- $\alpha$ .

Treatment of periodontitis with diabetes mellitus did not only include local factors, but also systemic factors handling blood sugar.7 such Researchers showed that the local periodontitis treatment such as scaling and root planing histologically not indicate new connective tissue attachment.<sup>5</sup> Periodontitis treatment of diabetes that has been done is the tissue surrounding dental treatments. antibiotics, and blood sugar control. While primary pathological the condition in people with diabetes



mellitus is a disorder of the healing process. 10,11

Sea cucumbers have long been used as a traditional medicine. One species of sea cucumbers which used as medicine is Stichopus hermanii that contains nutrients in wound healing, as an anticoagulant and antithrombotic, decrease cholesterol level and blood fat. anti-cancer, anti-tumor, antirheumatic. immunostimulant. antimalarials, antiviral, antifungal and antibacterial.<sup>12</sup> The content of omega-3 (EPA and DHA) in Stichopus hermanii can decrease cholesterol levels because infection of mix periodontopatogen bacteria and served to accelerate the healing of wounds and also as an antiinflammatory that inhibits production of TNF-α and IL-1β. 13,14 According to research of Rizal (2012)<sup>15</sup>, Stichopus hermanii is contained by flavonoids. The content of flavonoids as antioxidants that indirectly supporting anti-inflammatory effects by inhibiting the release of inflammatory mediators. 15,16

Sea cucumbers are one of the marine life that used in the health field. Sea cucumbers have shown a beneficial effect on periodontal therapy as well as patients with diabetes mellitus. In addition to cucumbers, one therapy that is emerging today is hyperbaric oxygen therapy (OHB).

Hyperbaric Oxygen Therapy (OHB) is a treatment in patients with giving 100% pure oxygen by inhalation method in hyperbaric hyperoxia at the air pressure of more than 1 atmosphere Absolute (ATA) and has been used as adjuvant therapy in various diseases.<sup>10</sup>

Hyperbaric Oxygen Therapy (OHB) shown a beneficial effect on periodontal tissues by increasing oxygen pressure in the pocket. In periodontitis

patients with diabetes mellitus who use OHB therapy there is decreased levels of malondialdehyde (MDA), which is one biomarker of oxidative stress. <sup>17,18</sup> In addition, OHB therapeutic effects on monocytes and macrophages are causing disruptions to production of proinflammatory cytokines, resulting in a reduction in pro-inflammatory cytokines in conditions of oxidative stress. <sup>19</sup>

The research results of Fitria et al (2013),<sup>20</sup> cucumbers therapy for 7 days effectively decrease blood sugar levels as well as the golden sea cucumber is used as an antibacterial and antifungal in inhibiting the growth of bacteria. The research results of Prabowo et al (2014),<sup>21</sup> OHB therapy 100% 2.4 ATA 3 x 30 minutes a day with a pause of 5 minutes for 7 days decreased blood sugar levels in diabetes mellitus type 2. It is necessary to investigate the combination of sea cucumber powder gold and OHB therapy against TNF-α expression in diabetes mellitus induced P. Gingivalis bacteria.

#### MATERIALS AND METHODS

The research is true experimental research with factorial design study. Parameters were examined in this study was the expression of TNF  $\alpha$  in diabetes mellitus and periodontitis. The samples were 20 Wistar rats (Rattus novergicus Wistar strain) were divided into five groups, where the selected criteria is the type males, aged 3-4 months with a body weight of 150-200 grams.

The tools used in this research is the syringe 5 cc and syringes 3cc, syringes insulin 1 mL, blood glucose test strips, gauges blood sugar, animal cages, food and drink rat, scales,



sodium carboxymethylcellulose (CMCNa) 2%.<sup>22</sup>

ISSN: 1907-5987

measuring cups and a stirrer, test tubes, test tube rack, micropipette, micro brush, glass, glass slide, surgical scissors, microscope, chamber of animals, immunohistochemistry kit.

Materials used nicotinamide 230 mg/kg, 200 mg canamycin, 200 mg ampicillin, 0.6% chlorhexidine gluconate, streptozotocin (to make the rats diabetes mellitus), 10% dextrose, citric acid buffer 0.05M pH 4.5, aluminum foil, bacteria Porphyromonas gingivalis ATCC 33277, powder golden sea cucumber (Stichopus hermanii), phosphate buffered saline (PBS), 4.13% EDTA, xylol, TNF-α antibody (sc-52 746-Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA ), H2O2, aquadest, Wistar rats, Wistar rats standard food, beverages Wistar rats (ordinary tap water), 100% pure oxygen in a hyperbaric chamber.

The division of research subjects. Those wistar rats were divided into five group: first, negative control group (K0); second, groups of diabetic periodontitis rats (K1); third, group of diabetic periodontitis rats with treatment golden sea cucumber (K2); fourth, groups of diabetic periodontitis rats with treatment of hyperbaric oxygen (K3); and five, groups of diabetic periodontitis rats with treatment combination golden sea cucumber and hyperbaric oxygen (K4).

Making the golden sea cucumber powder. Dissected sea cucumber to remove all viscoeral organs, then blowdry sea cucumber using freeze dryer at a temperature of 2-8 ° C with a pressure of 5 mTorr. The dried sea cucumbers are weighed and then blended into a powder. Making the golden sea cucumber powder gel using a mixture of

Diabetes Induction procedures for Wistar rats. Day 7 after adaptation, induction of streptozotocin (STZ). DM induction done by is giving nicotinamide (NAD) of approximately 230 mg / kg, dissolved in the liquid PBS 10 minutes and then given STZ intraperitoneal for 15 minutes injected a single dose of 65 mg/kg in rat that had fasted overnight between 8-12 hours. Then rats were incubated for 7 days. Otherwise DM rats when blood glucose levels over 220 mg / dl.<sup>23</sup>

**Procedure Premedication before induction of bacteria**. All the rats were given canamycin (20 mg) and ampicillin (20 mg) daily for 4 days in drinking water, and oral cavity swabbed with 0.12% chlorhexidine gluconate as antiseptic.<sup>24</sup>

**Procedure Preparation and Induction Bacteria.** Induction of bacteria using ATCC P.gingivalis imposing 33277 with 2 ml of 1 x 109 cells / ml of live bacteria in PBS, and peroral administration. Most bacteria applied using cotton bud along the gingival sulcus and inserted of anal to the colorectal region with syringe canula. Then incubation for 3 weeks since giving bacteria counted first. <sup>25</sup>

After the rats in the induction of STZ and P.gingivalis bacteria, rats were given a therapeutic treatment. For K2 and K4 given group therapy treatment gel golden sea cucumber powder 3% applied topical as much as 0.1 mg / day for 7 days using microbrush on inflamed sulcus gingival. K3 and K4 group treated OHB therapy, administration of 100% oxygen 2.4 ATA 3 x 30 minutes a



day with a pause of 5 minutes for 7 days. 26.21

On day 8 after therapy gel golden sea cucumber and OHB, all groups were sacrificed then performed euthanasia by the neck (cervical) dislocation, the dosage is taken on the part of the mandible. After that. the decalcification phase part of the mandible for 4 weeks using 4.13% EDTA. Furthermore. the processing and staining by immunohistochemistry (IHC) was performed using the streptavidin-biotinperoxidase labeled streptavidin-biotin (Dako, Carpinteria, USA), and then observed using a microscope 1000x magnification and calculated as much as 20x field view.

Data were analyzed using parametric statistical test one way ANOVA, with a test for normality using the Shapiro-Wilk test, then continued homogeneity statistic test using Levene test. Then the calculation results


compared (comparing the expression of TNF- $\alpha$  in each treatment group).

### **RESULT**

Before doing hypothesis test between groups, first of all each group tested the normality by using statistical test of Shapiro-Wilk, because in this study sample size <50 and the results obtained is Shapiro-Wilk test has a significant value of p> 0.05, so that the distribution of data on this research is abnormal.


**Table 1.** The average and expression deviation standard of TNF- $\alpha$  in each group

Group	Average ± Deviation Std.
<b>K</b> 0	$5.00 \pm 0.816$
<b>K1</b>	$11.50 \pm 1.291$
<b>K2</b>	$2.50 \pm 0.577$
<b>K3</b>	$8.25 \pm 2.217$
<b>K4</b>	$3.00 \pm 0.816$



**Figure 1.** Average and Deviation Standard of TNF-α expression in each group





**Figure 2.** TNF- $\alpha$  expression in macrophages in the periodontal ligament rat (Rattus norvegicus) with 400x magnification. A. K0 (normal); B. K1 (diabetes + bacteria); B. K2 (diabetes + bacteria + sea cucumbers); D. K3 (diabetes + bacteria + OHB); E. K4 (diabetes + bacteria + OHB + sea cucumber). Expression of TNF- $\alpha$  are marked with brown color in macrophages (arrows).

It was found that the expression of TNF- $\alpha$  have variances were not homogenous because p=0.041 (p<0.05). Therefore, it can be concluded that there is a difference of variance between sets of data are compared.

The research data were not normally distributed, and found that the data are not homogeneous, then doing non parametric Kruskal Wallis hypothesis test to determine the expression of TNF- $\alpha$ .

Table 2. Kruskal Wallis test Result

	Sig.
TNF-α expression	0.002

In the table 2. obtained significance value of 0.002 (p <0.05). It shows a significant difference in each group under these conditions, followed by Mann-Whitney test to find out where is the difference in each group with a significance level P <0.05.

Tabel 3. Mann-Whitney test Result

	K0	<b>K</b> 1	<b>K2</b>	К3	<b>K4</b>
K0		0.020*	0.019*	0.028*	0.027*
<b>K</b> 1			0.019*	0.059	0.020*
<b>K2</b>				0.019*	0.343
К3					0.020*

ISSN: 1907-5987

## **DISCUSION**

Nicotinamide (NAD) is an antioxidant which exerts protective effect on the cytotoxic action of STZ by scavenging free radicals and causes only minor damage to pancreatic beta cell mass producing type 2 diabetes.<sup>23</sup>

Streptozotocin is a drug that has a specific toxicity in pancreatic  $\beta$  cells. Streptozotcin cause DNA alkylation and  $\beta$  cell death because STZ enter cells through the glucose transporter GLUT2, toxic mechanism is not specific for  $\beta$ 

<sup>\*</sup>there is significant difference



cells and can cause damage to other including the liver kidneys.<sup>27,28</sup> Damage to the pancreatic beta cells causes the body can not produce insulin, causing blood glucose levels to rise (a state of hyperglycemia). Conditions of hyperglycemia can result in the formation of reactive oxygen species (ROS) which would then trigger an active nuclear factor kappa β (NFkβ) that play a role in the increased production pro-inflammatory of cytokines. <sup>23,29</sup> According to research from Swaroop et al (2012) 30 showed that TNF- $\alpha$  can play a potentially important pathophysiological role in the development of insulin resistance. Results of research on groups of bacterial-induced diabetic rats found increased expression of TNF-α compared with normal rats, it was shown that the group of bacterialinduced diabetic rats occurred diabetes.

Porphyromonas gingivalis is an anaerobic gram-negative rod bacteria and include black-pigmented bacteria, these bacteria cause inflammation and tissue damage in diseases periodontal.<sup>31,32</sup> P.gingivalis get into crevices using fimbriae and attached digingiva, epithelial cells gingival proteases have the ability to destroy periodontal tissue directly or indirectly, cause inflammatory and LPS can responses and resulting in production of cytokines in periodontal tissue, cause the activation of genes and cause bone damage alveolar. 31,6 Based on the results of the research showed that there were significant differences (p=0.020)TNF-α expression on between groups of normal rats and a group of bacterial-induced diabetic rats (see Table 5). The test results showed that the expression of TNF- $\alpha$  in the group of bacterial-induced diabetic rats

 $(11:50 \pm 1.291)$  increased compared with normal rats  $(5.00 \pm 0816)$ , this is due to streptozotocin-induced rats to make samples of rats became diabetic and bacteria capable P.gingivalis inducing the sample into periodontitis.

Some research it is known that the golden sea cucumber (Stichopus hermanii) as a potential therapy in the field of dentistry, besides the golden sea cucumber (Stichopus hermanii) itself has long been used as a traditional medicine. Golden sea cucumber powder used on research using the golden sea cucumber (Stichopus hermanii) which are found in waters of Indonesia.<sup>33</sup>

Based on the results of the research showed that there were significant differences on the expression of TNF-α between groups of normal groups of bacterial-induced rats, diabetic rats and rats treated groups of sea cucumbers (see table 5). The test results showed that the expression of TNF- $\alpha$  in groups of rats treated cucumbers decreased compared to the group of normal rats and a group of diabetic rats induced bacteria, because the content of the golden sea cucumber that is therapeutic as it helps wound healing, antibacterial, antifungal, antitumor, antianaphylactic, antiinflammatory, antinociceptive and antioxidant.<sup>34</sup>

(Stichopus Teripang gold hermanii) examined contains the active ingredient antibacterial. antifungal (antifungal), antitumor anticoagulant (anti-clotting) and it also contains compounds that can reduce cholesterol and lipid, anticancer and antitumor compounds, as well antibacterial compounds.35 Golden sea cucumber has a various kinds of content, one of the active ingredients of the golden sea cucumber is Triterpenoid



strong role as an immunostimulatory. In general Triterpenoid group capable of release of damaging the cell membrane, deactivate enzymes and proteins denature so that the cell wall is damaged due to a decrease in permeability. Changes intervals for 7 cytoplasmic permeability of the membrane allows ions important organic entry into the cell resulting in significantly. 21,40,26 poor growth and even to kill cells. 36.35 In addition, the gold content of sea cucumbers that PUFAs (DHA and EPA) CONCLUSSION may block the formation prostaglandins and proinflammatory

cytokines (IL-6, IL-1 $\beta$  and TNF- $\alpha$ ). <sup>37,38</sup> Hyperbaric oxygen increases the formation of oxygen free radicals, then oxidize proteins and lipid membranes of bacteria, destroys DNA and inhibits bacterial metabolic functions. enzyme superoxide dismutase, catalase, glutathione and glutathione reductase inhibiting free radical formation, until the oxygen concentration exceeded enzyme concentration.<sup>39</sup> Based on the results of the research showed that there were significant differences between groups of normal rats and rats treated group OHB, but there is no significant difference between groups of bacterialinduced diabetic rats and rats treated group OHB. The test results obtained by the expression of TNF-α in groups of rats treated group OHB increased compared to normal rats, but decreased the expression of TNF- $\alpha$  in groups of rats treated OHB compared bacterial-induced diabetic rats.

Based on the results of the study showed that there were significant differences between groups of bacterialinduced diabetic rats and mice combination treatment group due to the content of one of the golden sea cucumber are flavonoids that act as antioxidants and indirectly support the anti-inflammatory effects by inhibiting mediators inflamasi.16Penurunan expression TNFα at 2.4 ATA hyperbaric oxygen 100% oxygen 3x30 minutes with 5 minute days clinically recommended, because it can lower blood sugar levels, mRNA expression and activity of antioxidant enzymes

ISSN: 1907-5987

According to the research, it can be concluded that there is the influence of the golden sea cucumber powder 3% and hyperbaric oxygen therapy 3 x 30 minute interval of 5 minutes for 7 days on the expression of TNF-α in rats with mellitus induced diabetes by bacterium Porphyromonas gingivalis.

#### REFERENCE

- Malik G, Lehl G, Talwar M. 2011. Association Of Periodontitis With Diabetes Mellitus: A Review. Journal of Medical. 1 14-10. Available http://gmch.nic.in/journalgmch/Archives/5 %20Review%20article-
  - $\underline{\%20Periodontitis\%20with\%20Diabetes\%2}$ Omellitus.pdf. Diakses 7 April 2015.
- Akhila S and Malaiappan S. 2014. Periodontitis and Diabetes- A Two Way Connection. Journal of Dental and Medical Sciences e-ISSN: 2279-0853, p-ISSN: 2279-0861, 13 (1): 61-58.
- Petersen PE and Ogawa H. 2005. 3. Strengthening the Prevention of Periodontal Disease: The WHO Approach. Periodontal, 76(12): 2193-2187.
- Preshaw PM, Alba AL, Herrera D, Jepsen 4. S, Konstantinidis A, Makrilakis K, Taylor R. 2012. Periodontitis and Diabetes: A Two-Way Relationship. Diabetolodia. PMCID: PMC3228943, 55(1): 8-1.
- Newman MG, Takei HH, Takei HH, 5. Klokkevold P, Carranza F. 2012. Caranza's Clinical Periodontology, 11th



- ISSN: 1907-5987
- Philadelphia: WB Saunders Company. P. 43-41.
- Morran MP, Alexander LA, Slotterbeck BD, McInerney MF. 2009. Dysfunctional Innate Immune Responsiveness Porphyromonas Gingivalis Lipopolysaccharide in Diabetes. Oral Microbiol Immunology, 24: 339-331.
- Anil S, Varma SV, Preethanath RS, Anand 7. PS and Al Farraj A. 2012. The Emerging Concepts on The Impact of Periodontitis on Systemic Health. Periodontal Disease - A Clinician's Guide. ISBN: 978-953-307-818-2. P. 164-131.
- Jiang Z, Cui Y, Gao R, Lia Y, Fuc Z, 8. Zhang B, andGuan C. 2013. Study of TNF- $\alpha$ , IL-1 $\beta$  and LPS Levels in The Gingival Crevicular Fluid of A Rat Model of Diabetes Mellitus and Periodontitis, 34: 304-295.
- Engebretson S, Chertog R, Nichols A, Hey-Hadavi J, Celenti R and Grbic J. 2007. Plasma Levels of Tumour Necrosis Factorα in Patients With Chronic Periodontitis and Type 2 Diabetes. J Clin Periodontol, 34: 24-18.
- Mulawarmanti D. 2008. Profil cAMP, NF-10. κβ, HIF-1α, i-NOS, dan MMPs Jaringan Periodontal Tikus Wistar Pada Kondisi Hiperglikemia yang Dipapar Hiperoksia Hipebarik. Disertasi. Surabaya : Program Pascasarjana Universitas Airlangga 2008.
- Gurdol F, Cimsit M, Oner-Iyidogan Y, 11. Kocak H, Sengun S, and Yalcinkaya-Demirsoz S. 2010. Collagen Synthesis, Nitric Oxide and Asymmetric Dimethylarginine in Diabetic Subjects Undergoing Hyperbaric Oxygen Therapy. Physiol, 59: 429-423.
- Farouk AE, Ghouse FAH and Ridzwan BH. 2007. New Bacterial Species Isolated from Malaysian Sea Cucumbers with Optimized Secreted Antibacterial Activity. ISSN 1553-3468. American Journal Biochemistry and Biotechnology, 3(2): 65-60.
- Sari RP and Revianti S. 2011. The activity 13. of Stichopus hermanii extract on triglyceride serum level in periodontitis. Dental Jurnal Majalah Kedokteran Gigi, 44(2): 110-106.
- Calder 2006. 14. PC. Long-chain Polyunsaturated **Fatty** Acids and Inflammation. Scandinavian Journal of Food and Nutrition, 50(S2): 61-54.
- Rizal MB. 2012. Komposisi Senyawa 15. Organik dan Anorganik Teripang Pasir dan Teripang Emas yang Berperan dalam

- Proses Pulp Healing. Skripsi, Fakultas Kedokteran Gigi Universitas Hang Tuah Surabaya. Indonesia.
- Hidayati NA, Listyawati S and Setyawan 16. AD. 2008. Kandungan Kimia dan Uji Antiinflamasi Ekstrak Etanol Lantana camara L. padaTikus Putih (Rattus norvegicus) Jantan. ISSN: 0216-6887. Bioteknologi, 5(1): 17-10. Available from http://biosains.mipa.uns.ac.id/C/C0501/C05 0102.pdf. Diakses 2 April 2015.
- Nogueira-Filho GR, Rosa BT, and David-Neto JR. 2010. Effects of Hyperbaric 17. Oxygen Therapy on The Treatment of Severe Cases of Periodontitis. Undersea and Hyperbaric Medical Society, Inc: 37(2): 114-107.
- 18. Mulawarmanti D and Widyastuti. 2008. Effect of Oxygen Hyperbaric Therapy on Malondialdehyde Levels in Saliva of Periodontitis Patients With Type 2 Diabetes Mellitus. Dental Jurnal Majalah Kedokteran Gigi, 41(4): 156-151.
- 19. Thom SR. 2011. Hyperbaric Oxygen-Its Mechanism and Efficacy. PlastReconstr Surg, 127(1): 141s-131s. Available at http://www.ncbi.nlm.nih.gov/pmc/articles/P MC3058327/. Diakses 21 Maret 2015.
- 20. Fitriah, Theodorus dan Kamaluddin MT. 2013. Efek pemberian serbuk teripang (Stichopus Variegatus) jangka panjang terhadap kadar gula darah tikus Putih Jantan Galur Wistar Model Hiperglikemik. MKS, 45(1): 10-5.
- Prabowo Sulistiana, Nataatmadja Maria, 21. Poernomo Hadi Janto, Dikman Irmawati, Handajani Fitri, Tehupuring Sihning E. J., Soetarso1 Iswahyudi, Mohammad Survokusumo Guritno, Aulanni'am, Herawati Anita, West Malcolm. 2014. Hyperbaric Oxygen Treatment in a Diabetic Rat Model Is Associated with a Decrease in Blood Glucose, Regression of Organ Damage and Improvement in Wound Healing. Health, 1958-1950. 6, Availabe from http://dx.doi.org/10.4236/health.2014.6152 28
- Mulawarmanti D, Parisihni K, Wedarti YR. 22. 2014. Catalase Activitty of Sea Cucumber Extract (Stichopus hermanii) Periodontitis induced by Porphyromonas Gingivalis. Regional Oral Biology Sientific Meeting 2014. Central Library, Universitas Indonesia. P. 30-26.
- Srinivasan K and Ramarao P. 2007. Animal 23. models in type 2 diabetes research: An



- ISSN: 1907-5987
- overview. Indian J Med Res. 2007 March; 125: 472-451.
- 24. Kesavalu L, Sathishkumar Bakthavatchalu V, Matthews C, Dawson D, Steffen M, and Ebersole JL. 2007. Rat Model of Polymicrobial Infection, Immunity, and Alveolar Bone Resorption in Periodontal Disease. American Society for Microbiology. INFECTION AND IMMUNITY. 75(4): 1712-1704. doi:10.1128/IAI.00733-06.
- Praptiwi. 2008. Inokulasi Bakteri dan 25. Pemasangan Cincin Ligatur untuk Induksi Periodontitis padaTikus. Majalah Kedokteran Gigi, 15(1): 84-81.
- Harnanik T. 2008. Efek Oksigenasi 26. Hiperbarik Terhadap Peningkatan Aktivitas Antioksi dan pada Penderita Diabetes Mellitus Tipe 2. Tesis. Surabaya: Program Pasca Sarjana. Universitas Airlangga. P. 87.
- Lenzen S. 2007. The mechanism of 27. Alloxan and Streptozotocin induced Diabetes. Diabetologia (2008), 51: 226-216.
- Deeds MC, Anderson 28. JM, Armstrong AS, DA Gastineau, HJ Hiddinga, A Jahangir, NL Eberhardt, and YC Kudva. 2011. Single Dose Streptozotocin Induced Diabetes: Considerations for Study Design in Islet Transplantation Models. Lab Anim, 140.doi: 10.1258/la.2010.010090.
  - PMCID: PMC3917305. Available from http://www.ncbi.nlm.nih.gov/pmc/articles/P MC3917305/. Diakses 11 Januari 2016.
- Mealey BL and Thomas WO. 2006. AAP-29. Commissioned Review Diabetes Mellitus Periodontal Diseases. Journal Periodontol, 77 (8): 1303-1289. 10.1902/jop.2006.050459.
- Swaroop JJ, Rajarajeswari Dand J.N. 30. Naidu. 2012. Association of TNF-α with insulin resistance in type 2 diabetes mellitus. Indian J Med Res, 135: 130-27.
- Kimura S, Ohara-Nemoto Y, Shimoyama 31. Y, Ishikawa T and Sasaki M. 2012. Pathogenic Factors of P. gingivalis and the Host Defense Mechanisms. ISBN 978-953-307-924-0. P. 17-1.
- Hussain M, Stover Cordula M, and Dupont 32. A. 2015. P.Gingivalis in periondontal disease and atherosclerosis-sense of action for antimicrobial peptides and complement. Mini Review Aticle. 2015 February; 6(45): 5-1.
- Bordbar S, Anwar F, and Saari N. High-33. Value Components and Bioactives from

- Sea Cucumbers for Functional Foods-A Review.Mar. Drugs. 2011 October 10; 9: 1761-1805. doi:10.3390/md9101761. ISSN 1660-3397.
- Shahrulazua A, Samsudin AR, Iskandar 34. MA, Amran AS. 2013. The In-Vitro Effects of Sea Cucumber (Stichopus sp1) Extract on Human Osteoblast Cell Line. Malaysian Orthopaedic Journal, 7(1): 41-8. Doi:
  - http://dx.doi.org/10.5704.MOJ.1303.015.
- Roihanah S, Sukoso, Andayani S. 2012. 35. Aktivitas Antibakteri Ekstrak Teripang Holothuria sp. Terhadap Bakteri Vibrio harveyi Secara In vitro. Fakultas Perikanan dan Ilmu Kelautan, Universitas Brawijaya,
- Winarni D, Handono CD, Sugiharto. 2013. 36. Efek Imunostimulatori Beberapa Fraksi Teripang Lokal Phyllophorussp terhadap Histopatologi Limpa (Musmusculus) Yang Diinfeksi Mycobacterium tuberculosis. Available from
  - http://journal.unair.ac.id/filerPDF/Cipto%2 0Dwi%20Handono.PDF. Diakses 13 April
- Kordi MGH. 2010. A to Z Budidaya Biota 37. Akuatik untuk Pangan, Kosmetik dan Obatobatan. Yogyakarta: Lily Publisher. H. 18.
- Heba El Bialy, Ola MG, and Khaled SA. 38. 2011. Conversion of oil waste to valuable fatty acids using Oleaginous yeast. World J Biotechnol, Microbiol 27: 2791-98. Available from http://www.academia.edu/723079/Conversi on of oil waste to valuable fatty acids using\_Oleaginous\_yeast. Diakses November 2015.
- Wibowo A. 2015. Oksigen Hiperbarik: Terapi Penyembuhan Luka. Int. J. Unila, 5 (9): 128-124. Available from http://download.portalgaruda.org/article.ph p?article=328322&val=5503&title=Oksige n%20Hiperbarik:%20Terapi%20Percepatan %20Penyembuhan%20Luka. Diakses 12 Desember 2015.
  - Matsunami T, Sato Y, Yuki H, Satomi A, Haruka K, Takuya S, Masayoshi Y. 2011. Original Article Enhancement of reactive oxygen species and induction apoptosis in streptozotocin-induced diabetic hyperbaric oxygen exposure. Int J Clin Exp Pathol, 4(3): 266-255. Available from http://www.ijcep.com/IJCEP1101009. Diakses 14 Januari 2016.